

Admissibility of initial data in spherical collapse

H. M. Antia

Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India
email: antia@tifrvax.tifr.res.in

Abstract

Gravitational collapse of a spherically symmetric cloud has been extensively studied to investigate the nature of resulting singularity. However, there has been considerable debate about the admissibility of certain initial density distributions. Using the Newtonian limit of the equations governing collapse of a fluid with an equation of state $p = p(\rho)$ it is shown that the density distribution has to be even function of r in a spherically symmetric situation provided $dp/d\rho \neq 0$ even in comoving coordinates. We show that recent claim by Singh that the discrepancy pointed out earlier is due to their use of comoving coordinates is totally incorrect. It is surprising that he expects the use of comoving coordinates to make any difference in this matter. It is also argued that the strong curvature naked singularities in gravitational collapse of spherically symmetric dust do not violate the cosmic censorship hypothesis.

There has been considerable discussion about the nature of allowed initial data in spherically symmetric collapse. Unnikrishnan [1,2] has argued that density has to be an even function of radial distance when a realistic equation of state with pressure is used. On the other hand, Jhingan, Joshi and Singh [3] have claimed that odd terms in density cannot be ruled out. Antia [4] has pointed out that their claim is incorrect as they have not included the continuity equation in their analysis. Recently Singh [5] has claimed that this difference is because of their using the Lagrangian (or comoving) coordinates. In this paper we show that claim of Singh is totally incorrect and even in comoving coordinates the density has to be even function of radial distance when the matter obeys an equation of state of the form $P = P(\rho)$ with $dP/d\rho \neq 0$.

Following Singh [5] and using his notation, we expand

$$R(t, r) = r(a_0(t) + ra_1(t) + r^2a_2(t) + \dots) \quad (1)$$

However, in the comoving frame as used in [3] at $t = 0$ the scaling is chosen to ensure $R(0, r) = r$, which yields

$$a_0(0) = 1, \quad a_i(0) = 0 \quad (i > 0) \quad (2)$$

Now according to [5] the coefficient a_1 satisfies the differential equation

$$\frac{da_1}{dt} = \frac{a_1}{a_0} \frac{da_0}{dt} \quad (3)$$

With the initial condition given by Eq.(2) this equation will yield $a_1 = 0$ and hence the proof given by Antia [4] for the Eulerian equations can be continued to show that all the odd terms in expansion for density will vanish even in the Lagrangian case. Admittedly, the proof will be more involved in this case as compared to the Eulerian case considered in [4]. If the scaling $R(0, r) = r$ is not used then a_1 may be non-zero but in that case r will not represent radial distance.

This result can be more easily seen if the continuity equation is written in the usual form given by eq.(3) of [3], i.e.

$$\rho(t, r) = \frac{F'(r)}{R^2 R'} \quad (4)$$

where

$$F(r) = \int_0^r \rho(0, r) r^2 dr \quad (5)$$

Thus we can expand

$$F(r) = r^3(F_0 + rF_1 + r^2F_2 + \dots) \quad (6)$$

Further, since ρ_1 the linear term in the expansion of density vanishes it immediately follows that $F_1 = 0$. Now using these expansions in Eq.(4) it follows that $a_1(t) = 0$. Thus the mistake of Singh [5] is clear, although he has differentiated the continuity equation with respect to time t he has not used the initial conditions to obtain the solution. It is not clear why he has not used the form of continuity equation as used by them in [3]. As a result, the claims made by Singh [5] are totally incorrect.

It is surprising that Singh thinks that the Lagrangian form of equations will give different results as compared to the Eulerian form, since at least for $t = 0$, $R = r$ and one cannot have a different expansion in the two frames. Further, it is clear that the continuity equation was not used in [3], and the attempt by Singh [5] to use the continuity equation is merely a poor after thought.

As far as instability of strong curvature naked singularities in spherically symmetric dust collapse is concerned, Antia [4] has already argued that the proof is valid for the general case considered by Singh & Joshi [6]. In fact Jhingan, Joshi and Singh [3] have themselves shown that strong curvature naked singularities are non-generic and arise from initial data set of measure zero. If that is true such solutions are clearly unstable, since there would be some allowed perturbation which will not be in the class of solutions leading to strong curvature naked singularity. Further, Singh [7] has stated the cosmic censorship hypothesis as ‘‘Gravitational collapse of physically reasonable matter starting from generic initial data leads to the formation of a black-hole, not a naked singularity’’. Thus clearly according to Singh, the counterexamples with strong curvature naked singularities [3,6,8] in gravitational collapse of spherically symmetric dust do not even violate the cosmic censorship hypothesis, because these do not arise from generic initial data!

Acknowledgments: I would like to thank Professor Pankaj S. Joshi, Dr. T. P. Singh and Dr. I. H. Dwivedi for a series of long and stormy discussions.

- [1] C. S. Unnikrishnan, Gen. Rel. Grav. **26** (1994) 655.
- [2] C. S. Unnikrishnan, Phys. Rev. **D53** (1996) R580.
- [3] S. Jhingan, P. S. Joshi and T. P. Singh, Class. Quantum Grav. **13** (1996) 3057.
- [4] H. M. Antia, gr-qc/9701023.
- [5] T. P. Singh, gr-qc/9702023.
- [6] T. P. Singh and P. S. Joshi, Class. Quantum Grav. **13** (1996) 559.
- [7] T. P. Singh, gr-qc/9606016.
- [8] P. S. Joshi and T. P. Singh, Phys. Rev. **D51** (1995) 6778.